

Regulation of phosphate transport in proximal tubules

J. Biber · N. Hernando · I. Forster · H. Murer

Received: 30 May 2008 / Accepted: 13 August 2008 / Published online: 29 August 2008
© Springer-Verlag 2008

Abstract Homeostasis of inorganic phosphate (P_i) is primarily an affair of the kidneys. Reabsorption of the bulk of filtered P_i occurs along the renal proximal tubule and is initiated by apically localized Na^+ -dependent P_i cotransporters. Tubular P_i reabsorption and therefore renal excretion of P_i is controlled by a number of hormones, including phosphatonins, and metabolic factors. In most cases, regulation of P_i reabsorption is achieved by changing the apical abundance of Na^+/P_i cotransporters. The regulatory mechanisms involve various signaling pathways and a number of proteins that interact with Na^+/P_i cotransporters.

Keywords Sodium-dependent cotransport · Phosphate homeostasis · Regulation · Phosphate excretion

Introduction

If one considers the diversity of biological functions of inorganic phosphate (P_i), it should come as no surprise that an organism requires mechanisms to maintain the extracellular P_i reasonably constant. Deviations of the extracellular concentration of P_i will impair a variety of physiological processes such as bone metabolism, cellular energetics, protein synthesis, and signaling cascades. In humans, both hypo- and hyperphosphatemia have severe clinical con-

sequences that underscore the need for stable and well-managed P_i homeostasis [51, 62].

In mammals, including humans, plasma concentration of P_i is determined by the intestinal intake of P_i , the excretion of P_i via the feces, the release of P_i from bone and soft tissue, and the renal excretion of P_i . It is the latter mechanism that is of greatest importance for whole body homeostasis of P_i and therefore needs to be tightly controlled. In quantitative terms, human kidneys filter approximately 200 mmol (or ~20 g) of P_i per day. In the steady state, approximately 15% or 30 mmol (~3 g) P_i of the filtered load is excreted by the kidneys. Under physiological equilibrium conditions, the total daily renal and fecal P_i excretion is approximately equal to the amount of intestinally absorbed P_i for a normal diet. However, renal excretion of P_i does not always operate at a constant rate, but it is permanently adjusted according to the daily fluctuations of numerous hormones, including the recently described phosphatonins, and metabolic factors [11, 108, 12] (Table 1). This implies that the transporters involved in reabsorbing P_i in the proximal tubule are subject to regulatory pathways that (in most cases) affect the abundance of apical Na^+/P_i cotransporters [84, 108]. Moreover, growth and age as factors affecting renal P_i handling must also be considered as potential regulators of P_i transporters [110].

This review focuses on those Na^+/P_i cotransporters localized in the apical membrane of proximal tubules that are involved in the renal P_i reabsorption and which respond to P_i homeostatic regulatory factors. In this context, we will concentrate on the mechanisms that regulate P_i excretion via modification of apical expression of Na^+/P_i cotransporters under both normal physiological and pathophysiological conditions. As most of the data comes from studies on rodents, one should exercise caution when extrapolating

J. Biber · N. Hernando · I. Forster · H. Murer
Institute of Physiology and Zurich Center for Integrative
Human Physiology, University of Zurich,
Zurich, Switzerland

J. Biber (✉)
Institute of Physiology, University of Zurich,
Winterthurerstrasse 190,
CH-8057 Zurich, Switzerland
e-mail: JuergBiber@access.uzh.ch

Table 1 Factors affecting proximal tubular phosphate reabsorption

Reabsorption decreased by	Reabsorption increased by
Parathyroid hormone	Growth hormone
Dopamine	Insulin-like growth factor
Phosphatonins (FGF-23; sFRP-4; MEPE)	1,25(OH) ₂ D3
Glucocorticoids	Phosphate depletion
Atrial Natriuretic Peptide	
Phosphate loading	
Metabolic acidosis	
Carbonic anhydrase inhibitors	
Estrogen	

the conclusions to higher mammals, including humans. For more integrated views of whole body P_i homeostasis and specific details of genetic and acquired diseases that impair renal P_i handling, the reader should refer to the following articles [14, 17, 55, 70, 91, 108, 113, 117, 126].

Localization and composition of the renal machinery for P_i reabsorption

The P_i concentration in the glomerular ultrafiltrate is approximately equal to that in the plasma. Under normal physiological conditions, approximately 80% of P_i contained in the primary urine is reabsorbed unidirectionally along the proximal tubules. The amount of P_i reabsorbed in proximal convoluted tubules is up to threefold higher than that in proximal straight tubules. There is no evidence for reabsorption of P_i along the loop of Henle, but some P_i may be absorbed along distal tubular segments; however, the identity of the distal tubular transporters is unknown. Moreover, it is unclear if P_i handling in the distal nephron is relevant for the systemic control of P_i [for review see, 11, 108].

Uptake of P_i at the brush border membrane of proximal tubular cells is strictly dependent on the presence of Na^+ ions [121], which indicates that there is no paracellular transport of P_i via cell–cell contact sites. Indeed, all P_i transporter proteins identified so far and that are localized at the brush border membrane of proximal tubule cells are Na^+ -dependent. They mediate secondary-active transport by coupling to the free energy of the transmembrane Na^+ gradient that is maintained by the Na^+/K^+ -ATPase localized at the basolateral membrane (Fig. 1). In theory, this would allow sufficient accumulation of P_i within the cell to provide the driving force for the exit step(s) of P_i at the basolateral side. No detailed information is currently available about the intracellular concentration of free P_i or the molecular identity of P_i transporters localized at the basolateral membrane.

It has been established that Na^+/P_i cotransport through the brush border membrane is mediated chiefly by gene products of the solute carrier family SLC34 [83]; designated SLC34A1 (NaPi-IIa) and SLC34A3 (NaPi-IIc). Both SLC34A1 and SLC34A3 are localized at the brush border membrane of proximal tubule cells and have not been detected in other segments of the nephron [32, 99]. Under normal dietary conditions, these proteins are most abundant in S1 segments of both cortical and juxtamedullary nephrons, with the highest abundance in the latter. In contrast to NaPi-IIa, which gradually decreases along the entire length of the proximal tubule, NaPi-IIc has not been detected in the S3 segments of rat and mouse kidneys (Fig. 1) [32, 90].

A gene product of the solute carrier family 17 (SLC17A1 or NaPi-1) was identified based on Na^+/P_i cotransport in an expression cloning study using *Xenopus laevis* oocytes and was also localized to the brush border membranes of proximal tubules [33]. However, follow-up studies indicated that this cotransporter is most likely not involved in proximal tubular reabsorption of P_i , but rather in the transport of organic ions [21].

Finally, recent findings indicate that a gene product of the solute carrier family 20 (SLC20A2 or Pit-2) [29]) should also be considered as a potential candidate for proximal tubular P_i reabsorption, as this protein was localized exclusively at the brush border membrane of




Fig. 1 The renal P_i -reabsorption machinery in rat and mouse proximal tubules. Na^+/P_i -cotransporters NaPi-IIa, NaPi-IIc and Pit-2 are localized at the brush borders of proximal tubules. NaPi-IIa and Pit-2 proteins are detected in S1, S2 and S3 segments, whereas NaPi-IIc cotransporter is absent in S3 segments

proximal tubules (Villa-Bellosta R, Ravera S, et al., unpublished observations). Its contribution to proximal tubule P_i reabsorption and significance for whole body P_i homeostasis remain to be determined.

How do different Na^+/P_i cotransporters contribute to renal P_i reabsorption?

Recent studies indicate that to answer this question, we must take account of the species and developmental stage of the animals under investigation. Furthermore, the finding that NaPi-IIa, NaPi-IIc, and Pit-2 show different time courses of dietary or hormonal regulation suggests that the underlying signaling pathways differ.

Knockout of the gene encoding NaPi-IIa (Npt2) established that in adult mice, P_i reabsorption is mediated largely by NaPi-IIa [7]. In brush border membrane vesicles (BBMV's) isolated from Npt2^{-/-} mice, Na^+ -dependent P_i uptake was reduced by 70%, which could account for the observed hypophosphatemic state of these animals. The remaining 30% of Na^+/P_i cotransport in BBMV's has been attributed to the activity of NaPi-IIc as its abundance increased in Npt2^{-/-} mice [115]. Taken together, these observations suggested that in adult wild-type mice, Na^+/P_i cotransport via NaPi-IIc plays only a marginal role. In agreement, NaPi-IIc abundance was found to be highest in weaning mice and decreased with age [99]. This conclusion is further supported by the lack of a phenotype in NaPi-IIc^{-/-} mice [79].

In humans, particularly adults, the contribution of the NaPi-IIc to renal reabsorption of P_i and to whole body P_i homeostasis appears to be more important than in mice. Mutations in the SLC34A3 gene have been described in patients with hereditary hypophosphatemic rickets with hypercalcuria (HHRH) by two independent studies [9, 75]. The renal wasting of P_i associated with this disease can therefore be correlated with an impaired function of NaPi-IIc. This suggests that in adult humans, NaPi-IIc contributes significantly to renal P_i reabsorption under normal steady-state conditions.

On the basis of in situ hybridization, expression of both members of the SLC20 family [29], Pit-1 and Pit-2, was detected along the entire nephron [116]. As noted above, the Pit-2 protein was recently localized in the brush border membranes of proximal tubules (Villa-Bellosta R, Ravera S et al., unpublished observations), but its contribution to renal P_i reabsorption remains unknown.

Na^+/P_i cotransporters at work

The transport kinetics of NaPi-IIa, NaPi-IIc, and Pit-1, -2 have been characterized in detail using the *Xenopus laevis*

oocyte expression system [for review see 38, 124]. All four proteins prefer Na^+ ions as the driving substrate, and P_i as the driven substrate, yet there are important differences that could influence their respective physiological functions.

First, NaPi-IIa is electrogenic, whereas NaPi-IIc is electroneutral. This finding is consistent with their different transport stoichiometries. Both prefer divalent P_i (HPO_4^{2-}); NaPi-IIa displays a $Na^+:P_i$ stoichiometry of 3:1, whereas for NaPi-IIc, the stoichiometry is 2:1. Thus, the electrogenicity of NaPi-IIa arises from the translocation of an additional Na^+ ion, at greater energetic cost to the cell. Despite this fundamental difference, at neutral pH, both NaPi-IIa and NaPi-IIc show similar apparent affinities for Na^+ (~50 mM) and P_i (<0.1 mM) as well as a qualitatively similar decreased transport activity with external acidification [38, 124, 125].

Second, in contrast to NaPi-IIa/c, Pit-1/2 prefer monovalent P_i ($H_2PO_4^-$). Like NaPi-IIa, both Pit-1 and Pit-2 are electrogenic, yet Pit-1 has been shown experimentally to display a 2:1 $Na^+:P_i$ stoichiometry. Apparent affinities for Na^+ are in the range of 50 to 70 mM and for P_i ~0.1 mM [18, 93, 124]. Unlike NaPi-IIa/c, Li^+ can also support P_i translocation by Pit-1, but at a significantly reduced rate and there is a weak affinity for arsenate [93].

Third, the pH dependence of NaPi-IIa and NaPi-IIc is similar; transport rates approximately double between pH 6.5 and 8 [124]. Such a pH-dependence may have to be considered when assessing renal excretion of P_i under conditions of an altered acid base status of the body [46, 87]. In contrast, the predicted maximum transport velocity for Pit-1/2-mediated Na^+/P_i cotransport is quite constant within this pH range [93].

Finally, until recently, phosphonoformic acid (PFA) was considered a general competitive inhibitor of Na^+/P_i cotransport [112]. Current data indicate that it inhibits only Na^+/P_i cotransport mediated by SLC34 family gene products. The K_i for PFA is approximately tenfold higher for Pit-1, -2 compared to that reported for NaPi-IIa. This may be a direct consequence of the different preference for P_i of the two families as PFA is expected to be divalent under normal physiological conditions [93].

Currently, no information about the three-dimensional structure of Na^+/P_i cotransporters is available. Based on extensive studies using cysteine scanning combined with functional studies and in vitro transcription/translation approaches, SLC34 proteins are predicted to comprise eight transmembrane domains with intracellular N- and C-termini. A large extracellular loop region links two halves of the protein that contain conserved repeated sequences also found in bacterial homologs. The repeats are proposed to form opposed reentrant loops that most likely constitute the transport pathway [38, 92, 124]. The secondary structure of SLC20 proteins is less well defined: they are

predicted to comprise ten transmembrane domains with extracellular N- and C-termini, however evidence for an obvious repeat region is so far lacking, and there is no homology with the SLC34 gene products in the proposed substrate translocation region [18, 98].

The functional unit of NaPi-IIa is a monomer [63]. Interestingly, a split-ubiquitin yeast two-hybrid assay, and subsequent studies provided evidence that NaPi-IIa may homodimerize [42]. Although apparently not relevant for transport function, one can speculate that homotypic interactions of NaPi-IIa proteins may be of relevance for the regulation of this cotransporter.

Apical positioning of Na⁺/P_i cotransporters

After correct targeting and insertion into the plasma membrane of epithelial cells, specific protein–protein interactions may be required to stabilize the final localization of membrane proteins [130]. Classical yeast two-hybrid screens performed against the C-terminus of NaPi-IIa

revealed interactions of this cotransporter with several PDZ domain (PSD-95, discs-large, ZO-1) containing proteins that may contribute to the stabilization of NaPi-IIa at the apical membrane (Fig. 2b) [15, 42–44]. Robust interactions of NaPi-IIa with single PDZ domains of the sodium–hydrogen exchanger regulatory factor (NHERF), protein family [104], NHERF1 and 2, NHERF3 (PDZK1) and NHERF4 (PDZK2) have been demonstrated in in vitro studies. Members of the NHERF family were proposed to scaffold several other membrane proteins, including transporters and receptors as well [43, 44]. Furthermore, NHERF1/2 bind to members of the merlin–ezrin–radixin–moesin (MERM) protein family, thereby providing a link to the cytoskeleton [104].

The interactions of NaPi-IIa with PDZ domains of the NHERF family were shown to be primarily via a class I PDZ binding motif (TRL) located at the C-terminus [43, 44]. In opossum kidney (OK) cells, the TRL-motif was shown to be required for apical expression [49]. The roles of some of these interactions of NaPi-IIa with PDZ proteins have been elucidated using several mouse knockout




Fig. 2 Regulation of the NaPi-IIa Na⁺/P_i-cotransporter. **a** Activation of protein kinase dependent signaling pathways, e.g., by PTH or dopamine, provoke internalization of NaPi-IIa cotransporters at the intermicrovillar clefts. Internalized NaPi-IIa proteins are degraded in the lysosomes. **b** Schematic illustration of the apical sorting and positioning of NaPi-IIa by PDZ proteins [15, 43, 44, 104]. The

recently suggested role of Gabarap in determining apical abundance of NaPi-IIa [94] is also indicated. **c** Modulation of the interaction of NaPi-IIa with NHERF1 by protein kinases activated by PTH. Phosphorylation of NHERF1 [128] weakens the apparent affinity for NaPi-IIa, which is subsequently delivered to the endocytic machinery

models. NHERF1^{-/-} mice exhibited phosphaturia that could be explained by a decrease of the apical abundance of NaPi-IIa [105]. These findings indicate a stabilization effect of NHERF1 on the membrane expression of NaPi-IIa. In contrast, in NHERF2^{-/-} mice, urinary excretion of P_i and apical expression of NaPi-IIa were unaffected under normal physiological conditions [31]. Despite the robust in vitro interactions of NaPi-IIa with NHERF3, neither urinary excretion of P_i nor the abundance of NaPi-IIa was impaired in NHERF3^{-/-} mice [23].

In addition to the NHERF proteins, NaPi-IIa also interacts with the PDZ domain of Shank2E, a splice form of Shank2 [78]. Shank2E is localized at the brush border membrane of proximal tubule cells, and there is evidence that it may be involved in the endocytic machinery that determines the surface abundance of NaPi-IIa.

A recent study revealed a role for the GABA_A-receptor associated protein (Gabarap) in determining the apical amount of NaPi-IIa [94]. In BBMVs isolated from Gabarap^{-/-} mice, the amount of NaPi-IIa was increased, whereas NaPi-IIc was unaffected. In agreement with these findings, Na⁺/P_i cotransport was higher in BBMVs from Gabarap^{-/-} animals, which was reflected in a reduced urinary P_i excretion. The molecular mechanism underlying the action of Gabarap is unknown. Gabarap may exert its effect on the apical sorting as it interacts with the N-ethylmaleimide-sensitive factor NSF [61]. Alternatively, NHERF1 which was also up-regulated in Gabarap^{-/-} mice [94], may increase the stabilization of NaPi-IIa in the brush border membrane of these animals.

Currently, there is little information about proteins interacting with NaPi-IIc or Pit-2. NaPi-IIc was reported to interact with NHERF1 and NHERF3 [123], the consequences of these interactions for apical positioning and regulation of NaPi-IIc remain to be clarified.

Regulation of proximal tubular Na⁺/P_i cotransport

General considerations

Theoretical considerations predict that at equilibrium, secondary active Na⁺/P_i cotransport across the brush border membrane of proximal tubules would give rise to a concentrating capacity of 100:1 for NaPi-IIc and 10,000:1 for NaPi-IIa [45, 124]. Although precise values for the intracellular concentration of P_i are unavailable, it is estimated to be in the lower millimolar range [14]. Thus, for a given membrane protein density, the capacity of proximal epithelia to reabsorb P_i is not determined by the driving force but by the turnover rate of the transport cycle. Alteration of P_i reabsorption therefore could be achieved either by changes in the abundance of Na⁺/P_i cotransport-

ers, the apparent affinities for the substrates or the lipid composition, which could influence the preferred conformational state of the protein and modulate partial reactions that constitute the transport cycle.

Most of the known effects of hormones and metabolic factors (Table 1) on renal P_i absorption take place via alterations of the abundance of Na⁺/P_i cotransporters within the proximal brush border membrane. The membrane abundance of a Na⁺/P_i cotransporter depends on the rate of protein synthesis in the biosynthetic pathway, the rate of insertion into the apical membrane and the rate of endocytotic processes. Currently, we know more about down-regulation (via endocytosis) of Na⁺/P_i cotransporters (in particular NaPi-IIa) than about the possible regulation of the apical sorting and insertion of newly synthesized cotransporters into the brush border membrane. In the following section, we will present an overall picture of the current knowledge about the acute hormonal regulation of Na⁺/P_i cotransporters, such as, e.g., by parathyroid hormone. Other important regulators of apical Na⁺/P_i cotransporter abundance, such as phosphatonins, P_i diet and steroids are discussed in separate sections. Finally, a few cases in which changes of renal excretion of P_i do not match alterations of apical abundance of Na⁺/P_i cotransporters will be discussed.

Changes in renal P_i excretion related to alterations of Na⁺/P_i cotransporter abundance

Parathyroid hormone

The paradigm for the alteration of the apical abundance of Na⁺/P_i cotransporters is represented by the regulation of the NaPi-IIa protein by parathyroid hormone (PTH) (Fig. 2a). Upon stimulation of PTH receptors, a decrease of apical NaPi-IIa abundance occurs within minutes, without changes in mRNA levels [4, 58]. In contrast, several hours are required for a PTH-induced redistribution of NaPi-IIc [90, 103]. This indicates that (at least in rodents) acute adjustment of renal P_i reabsorption by PTH occurs mostly by a redistribution of NaPi-IIa. So far, no regulation of Pit-2 by PTH in proximal tubules has been described.

Stimulation of PTH receptors leads to a rapid internalization of NaPi-IIa via an endocytotic machinery located at the intermicrovillar clefts that was not associated with fluid phase uptake, but was shown to be mechanistically similar to a receptor-mediated process that involves clathrin-coated vesicles [4]. Although putative endocytic motifs such as tyrosine- or di-leucine-based motifs are contained within the amino acid sequence of NaPi-IIa, mutations of these motifs did not affect the retrieval of NaPi-IIa [50]. Instead, a dibasic amino acid motif (KR), most likely localized within

an intracellular loop, was found to be required for internalization and degradation of NaPi-IIa [56]. The absence of this KR motif in the NaPi-IIc protein could explain the different endocytic rates of the NaPi-IIa and NaPi-IIc cotransporters after PTH stimulation. It is of interest that the peroxisomal farnesylated protein PEX19 was reported to interact with NaPi-IIa via the KR motif, yet the precise role of PEX19 in the trafficking of NaPi-IIa has not been elucidated [54].

Unlike megalin [28], internalized NaPi-IIa is not delivered to recycling subapical endosomes but, instead, is routed from the early endosomes to the lysosomes where it is degraded [4, 59, 89] (Fig. 2a). No lysosomal association of NaPi-IIc was detected after stimulation with PTH [90, 103], which again indicates that any intracellular trafficking of NaPi-IIc after hormonal stimulation differs from that of NaPi-IIa.

Several observations suggest that the final reaction(s) of the PTH-induced signaling cascades leading to NaPi-IIa endocytosis may affect the affinity of the cotransporter for its interacting proteins. The signaling cascades involved in the action of PTH have been well described. Receptors for PTH are localized both at the apical and the basolateral site of proximal tubular cells and differ with respect to their activation by different PTH fragments and formation of intracellular second messengers [41]. Stimulation of basolateral PTH receptors by 1–34 PTH (via Gs) provokes an increase of the intracellular cAMP concentration and activation of protein kinase A-dependent pathways, whereas stimulation of apical PTH receptors by 1–34 PTH or 3–34 PTH (via Gq) results in the hydrolysis of PIP₂ and activation of pathways involving protein kinase C activity. Experiments performed on isolated rat proximal tubules indicated that stimulation of both, apical and basolateral PTH receptors, elicits a down-regulation of NaPi-IIa [119]. Apical PTH receptors are part of a signaling complex scaffolded by NHERF1 that also includes phospholipase PLCB2 [77]. The importance of such a spatial arrangement for the hormonal regulation of the abundance of NaPi-IIa was demonstrated in experiments using kidney slices derived from NHERF1^{-/-} mice. The coupling of apical PTH receptors to the down-regulation of NaPi-IIa was completely abolished in the absence of NHERF1, whereas down-regulation of NaPi-IIa induced by a stimulation of basolateral PTH receptors (via cAMP) remained intact [22].

In addition to PKA and PKC pathways, regulation of the amount of NaPi-IIa by a pathway that involves protein kinase G should also be considered. Early studies documented that part of the renal response to the atrial natriuretic factor (ANF) includes changes of Na⁺/P_i cotransport [48]. Later, it was demonstrated that increases of cGMP concentrations induced by ANP or by nitric oxide (NO) resulted in a down-regulation of NaPi-IIa [3].

An involvement of MAPK (ERK1/2) kinases in the PTH action has been suggested from experiments performed with opossum kidney (OK) cells [67]. Moreover, in ex vivo studies, using mouse kidney slices, down-regulation of NaPi-IIa initiated by 1–34 PTH was partially prevented by the ERK1/2 inhibitor PD098059 [5]. Similarly, down-regulation of NaPi-IIa was partially inhibited by this blocker when PKC or PKG pathways were pharmacologically activated separately. In contrast, inhibition of ERK1/2 completely abolished endocytosis of NaPi-IIa in response to direct activation of PKA [5]. Based on these observations, we can envisage a central role of the MAPK pathway as a point where the different pathways (PKA, PKC, and PKG) may converge (Fig. 2a). Interestingly, inhibition of renal P_i reabsorption by FGF-23 (see below) was also reported to involve activation of the MAPK kinase pathway [129].

Is MAPK always required for down-regulation of NaPi-IIa? Investigations related to the phosphaturic effect of dopamine [39] demonstrated that dopamine elicits a down-regulation of NaPi-IIa via stimulation of apical dopamine receptors D1 [2]. This D1-receptor-mediated effect on NaPi-IIa abundance was completely prevented by a blocker of PKA (H89) but not by inhibiting ERK1/2 MAPK kinases, which indicates that cAMP dependent pathways relevant for down-regulation of NaPi-IIa may differ, depending on which receptor is activated (Fig. 2a).

The subcellular location of NHERF proteins remains unaffected after PTH treatment. The implication of different kinases in the signaling cascades described above suggest that the stability of NaPi-IIa within apical heteromultimeric protein complexes, such as the one scaffolded by NHERF1, may be altered by protein phosphorylation. Studies addressing this question revealed that NaPi-IIa was not phosphorylated either basally or upon stimulation by PTH [34]. However, increased phosphorylation of NHERF1 in response to PTH has been observed in murine kidney slices and suggested that phosphorylation of NHERF1 could induce a dissociation of the NaPi-IIa/NHERF1 complex [34]. Recently, a more detailed study demonstrated that after activation of PKC and PKA pathways, a serine residue (Ser77) within the first PDZ domain of NHERF1 was phosphorylated [128] (Fig. 2c). Phosphorylation of Ser77 upon activation of PKC appeared to occur more directly compared to the PKA effect. Because NaPi-IIa interacts with the first PDZ domain of NHERF1 [43, 44] and Ser77 is located within the predicted binding domain, phosphorylation of Ser77 is thought to induce a conformational change resulting in a lowered affinity of the NaPi-IIa/NHERF1 interaction. In this respect, it is interesting to note that truncation of the PDZ binding motif of the CFTR channel increased its diffusional mobility within the membrane [47]. Similarly, by promoting phosphorylation

of NHERF1, PTH (and possibly other factors) could induce an increase of the mobility of NaPi-IIa along the microvillus axis that may be required for internalization at the intermicrovillar clefts.

Phosphatonins

Tumor-induced osteomalacia (TIO), autosomal-dominant hypo-phosphatemic rickets (ADHR), recessive hyperphosphatemic rickets (ARHR) and X-linked hypophosphatemic rickets (XLH) are associated with phosphaturia and defects in bone metabolism [126]. Associated with these diseases, various new factors, termed phosphatonins, have been identified and were shown to be involved in the observed renal P_i wasting. So far, the list of phosphatonins comprises fibroblast growth factor 23 (FGF-23), secreted frizzled related protein-4 (sFRP-4), matrix extracellular phosphoglycoprotein (MEPE), and fibroblast growth factor 7 (FGF-7). The roles of phosphatonins in whole body P_i -homeostasis under pathophysiological conditions has recently been reviewed [12, 17, 55, 91, 117, 126]. However, apart from their roles in disease conditions, phosphatonins may exert regulatory functions on renal P_i reabsorption also under normal physiological conditions, whereby interference by other factors cannot be excluded [97, 106, 107, 109]. Overall, little is known about the signaling pathways of phosphatonins in proximal tubule cells that result in alterations of the abundance of Na^+/P_i cotransporters. Current knowledge about the cellular mechanisms is briefly discussed below.

Fibroblast growth factor FGF-23 FGF-23 was identified in patients with autosomal dominant hypophosphatemic rickets (ADHR) by positional cloning [30]. The phosphaturic effect observed at elevated serum levels of FGF-23 correlated with a decrease of the abundance of NaPi-IIa and NaPi-IIc [79, 80]. Moreover, FGF23^{-/-} mice display the converse behavior with increased P_i reabsorption and hyperphosphatemia [107]. In ex vivo studies, using isolated proximal tubules, FGF-23 was reported to reduce NaPi-IIa abundance [6].

FGF-23 likely activates one of the known c-splice isoforms of FGF receptors (FGFR). Some in vitro studies suggested the interaction of FGF-23 with FGFR's 1c, 3c, and 4c; however, the identity of the FGFR responsible for the action of FGF-23 in proximal tubules is currently unknown. FGF-23 receptors belong to the family of receptor tyrosine kinases, and its activation appears to be dependent on the presence of sugar chains such as those contained in heparin or klotho [65, 66, 74]. It is significant that klotho (see below) is required for the conversion of canonical FGFR's into functional FGF-23 receptors [122]. The importance of klotho for FGF-23 action was further suggested from observations made using a klotho^{-/-} mouse

model that develops a phenotype similar as FGF-23^{-/-} mice [102]. In klotho^{-/-} mice, the increase of serum P_i concentration could be correlated with increased abundances of NaPi-IIa and NaPi-IIc despite a massive increase of FGF-23 serum levels.

Studies in OK cells provided evidence that the signaling pathway activated by FGF-23 involves MAPK/ERK1-2 kinases [129]. Similarly as described for the PTH-induced signaling pathway (see above), the action of FGF-23 was blocked by inhibitors of ERK1/2. In addition, in isolated proximal tubules, FGF-23 was reported to provoke an increase of the production of prostaglandin PGE2 that was also suggested to occur via the MAPK/ERK pathway [111].

Secreted frizzled-related protein-4 Secreted frizzled-related protein-4 (sFRP-4) is overexpressed in TIO patients presenting renal P_i wasting [for review see 12]. Infusion of sFRP-4 into rodents was shown to provoke phosphaturia that could be correlated with a decrease of NaPi-IIa [10]. sFRP-4 is an antagonist of the Wnt pathway and, indeed, after infusion of sFRP-4, a decrease of the phosphorylated form of β -catenin was detected. However, the time course and the precise mechanism by which sFRP-4 leads to down-regulation of NaPi-IIa needs to be explored further.

Matrix extracellular phospho-glycoprotein and fibroblast growth factor 7 Overexpression of two other factors matrix extracellular phospho-glycoprotein (MEPE) and FGF-7, has been noted in tumors of TIO patients as well [for review see 12]. MEPE was also identified in tissue from a patient with oncogenic hypophosphatemic osteomalacia [95]. Conflicting results have been reported with respect to a direct action of MEPE on renal P_i handling. A recent study demonstrated that in rats, infusion of recombinant MEPE elicited an increase of P_i excretion [35]. Together with data obtained from cell culture experiments [96], it appears that MEPE indeed participates in the control of proximal P_i reabsorption. However, the precise mechanisms of the action of MEPE are not known. Also, further investigation will be required to establish if other factors regulating P_i homeostasis also interact with the renal action of MEPE.

Only minimal information is available at present about FGF-7 regarding its role in renal P_i wasting as observed in TIO patients. Based on studies performed with OK-cells, FGF-7 inhibits Na^+/P_i cotransport [25].

Klotho

In addition to interaction with FGF receptors (see above), klotho may exert direct effects on the abundance of Na^+/P_i cotransporters in proximal tubules. The extracellular domain of klotho, which is anchored by a single transmem-

brane segment, shows ~40% homology to β -glycosidase family-1 enzymes and thus may enzymatically alter sugar moieties of membrane proteins [118]. In kidneys, klotho is expressed in distal convoluted tubules [65]. Interestingly, the extracellular domain of klotho is shed from the membrane and is detected in the blood, which indicates that this domain may act as a humoral factor [53, 65, 85]. Evidence for a direct action of klotho on NaPi-IIa abundance was reported from in vitro experiments using microperfused proximal tubules and isolated BBMVs [52]. Most interestingly, incubation of BBMVs with klotho reduced Na^+/P_i cotransport activity, an effect most likely was due to a change in the glycosylation of NaPi-IIa itself. These observations suggest that modification of the glycosylation moieties of NaPi-IIa may affect its apical stability, similar to that described for the calcium channel TRPV5 [27].

Dietary P_i intake

It has long been known that changes in the dietary intake of P_i affect renal excretion of P_i . Interestingly, the capacity of renal P_i reabsorption adapts to altered intake of P_i within less than an hour (acute adaptation) and remains adjusted during prolonged intake of dietary P_i (chronic adaptation) [71, 73].

Adaptation to altered P_i diet is independent of PTH, vitamin D3, and growth hormones [19, 24, 71, 120]. However, analysis of the mechanisms involved in such dietary adaptation can be complicated by the involvement of other factors. For example, serum levels of FGF-23 were reported to be dependent on the state of phosphatemia [127]. Furthermore, studies performed with diabetic rats indicated that insulin may have a permissive effect on the adaptive response to low P_i diet as the increase of apical Na^+/P_i cotransport was abolished in these rats [1]. Also, lowered plasma levels of P_i may affect the rates of metabolic pathways of proximal tubular cells, such as for example of gluconeogenesis [64].

Within the first four hours after ingestion of a low P_i diet, brush border Na^+/P_i cotransport increases due to an increase of the amount of NaPi-IIa [26, 71, 73]. Rapid up-regulation induced by low P_i diet was also observed for Pit-2 but not for NaPi-IIc [79 and Ravera S et al., unpublished observations]. Whereas acute changes of NaPi-IIa abundance are independent of transcriptional regulation, controversial findings regarding the chronic regulation have been reported. In weaning mice, a chronic low P_i diet elevates the amount of NaPi-IIa mRNA in addition to protein abundance. This effect was ascribed to a P_i responsive element located within the promoter region of the Npt2 gene [60]. However, in adult mice fed chronically with a low P_i diet an increase of NaPi-IIc mRNA but not of NaPi-IIa mRNA was observed [76].

The signaling mechanisms involved in dietary regulation are not fully understood. The rapid increase of brush border Na^+/P_i cotransport was not blocked by cycloheximide, which suggests that de novo synthesis of Na^+/P_i cotransporters might not be necessary [73]. Yet, so far, there is no clear evidence for the existence of an intracellular, subapical pool of Na^+/P_i cotransporters that could explain the rapid adaptive phenomenon. Based on the observation that the fall in the serum level of P_i after ingestion of a low P_i diet precedes up-regulation of NaPi cotransport [73], one might hypothesize the existence of one or more sensing mechanism(s) for P_i that eventually may be located in the proximal tubular cells.

Renal excretion of P_i decreases after ingestion of a diet rich in P_i due to a reduction of apical Na^+/P_i cotransporter abundance. High P_i diet induces internalization of both NaPi-IIa and NaPi-IIc, yet by distinct mechanisms. Whereas the amount of NaPi-IIa in BBMVs was decreased after 2 h, reduction of NaPi-IIc was detectable only after 4 h. As found for the PTH action, internalized NaPi-IIa (by high P_i diet) is routed to the lysosomes, whereas NaPi-IIc is redistributed from the BBMVs to a subapical compartment and does not undergo lysosomal degradation [59, 71, 101].

Acute regulation of renal P_i excretion has recently been investigated after application of a P_i bolus directly into the duodenum of rats [13]. This maneuver elicited an increase of renal excretion of P_i already within 15 min. It is noteworthy that this response preceded changes in the plasma P_i , was independent of PTH and the phosphatonins FGF-23 and sFRP-4 and was not observed in animals with denervated kidneys. These data suggest that renal P_i excretion may be part of an intestinal–renal axis that may include a novel (humoral) factor originating from the duodenal mucosa.

Steroid hormones

The influence of steroid hormones on renal Na^+/P_i cotransporters has been studied under chronic conditions. Although chronically administered, steroids could act via genomic mechanisms, nongenomic effects cannot be excluded. Furthermore, it has to be considered that other factors may be involved in the action of a particular steroid.

Glucocorticoids In addition to other effects in proximal tubules, such as stimulation of the rate of gluconeogenesis, glucocorticoids are also involved in the regulation of the renal handling of P_i [36, 39, 108]. In kidneys from rats that were treated chronically with dexamethasone, the amount of NaPi-IIa protein and mRNA was decreased, in agreement with increased excretion of P_i [72]. Furthermore, an inverse correlation between Na^+/P_i -cotransport in BBMVs and glucosylceramide and sphingolipid content was reported in

these studies. It remains to be determined if glucocorticoids may affect, besides NaPi-IIa, other apical Na⁺/P_i cotransporters as well.

Estrogen Chronic treatment of ovariectomized rats with estrogen results in hypophosphatemia and hyperphosphaturia. Renal wasting of P_i observed under these experimental conditions has been explained by reduced Na⁺/P_i cotransport in BBMVs due to a decrease of NaPi-IIa abundance [8, 37]. Moreover, estrogen treatment was shown to decrease the abundance of NaPi-IIa mRNA, whereas NaPi-IIc mRNA was unaffected. It remains to be determined if the reported effect of estrogen on NaPi-IIa is a direct or indirect one. Although the effect on NaPi-IIa was reported to be independent of PTH [37], the reduction in NaPi-IIa abundance could result from an indirect action of estrogen on dopamine receptors and/or the metabolism of dopamine in proximal tubules [68].

1,25(OH)₂-Vitamin D₃ (VitD₃) Although the profound effects of VitD₃ on the metabolisms and homeostasis of calcium and phosphate are well described [20, 108], conflicting results about direct effects of VitD₃ on P_i reabsorption in proximal tubules have been reported.

In thyroparathyroidectomized (TPTX) rats, chronic administration of VitD₃ was reported to inhibit P_i reabsorption in proximal tubules [82]. Although interference by PTH was excluded in this study, more recent data suggest that VitD₃ action on renal P_i handling could be indirectly, involving altered serum levels of phosphatonins [109]. For example, in intact rats, repeated injections of VitD₃ resulted in an increase of serum levels of FGF-23 [106]. Furthermore, FGF-23 impairs the renal metabolism of VitD₃ by suppressing the expression of 25-hydroxyvitamin D 1 α -hydroxylase mRNA and by stimulation of the expression of 25-hydroxyvitamin D-24 hydroxylase mRNA [88, 107].

A VitD₃ responsive element has been identified in the promoter region of the human NaPi-IIa gene, suggesting that regulation of renal P_i reabsorption by VitD₃ may occur at the transcriptional level [114]. However, in VitD₃ receptor and 1, α -OHase^{-/-} mice, the abundance of the NaPi-IIa protein in BBMVs was reported to be similar as the one of wild-type mice [24, 100], which indicates that at least in mice, the NaPi-IIa abundance may not be controlled by VitD₃ dependent transcriptional mechanisms.

Changes in renal P_i excretion not related to alterations of Na⁺/P_i cotransporter abundance.

Apart from the hormonal and metabolic settings that result in a change of the abundance of Na⁺/P_i cotransporters as

discussed above, a few observations indicate that altered renal P_i excretion can occur also without parallel changes of the amount of apical Na/Pi cotransporters.

Potassium depletion

In rats depleted of potassium, P_i excretion was increased and was in agreement with a decreased Na⁺/P_i cotransport in isolated BBMVs. However, this change of Na⁺/P_i cotransport was not correlated with a reduced abundance of NaPi-IIa, but instead by increased amounts of NaPi-IIa. [131]. These contradictory findings were explained by an alteration of the lipid composition, (e.g., glucosylceramide), and altered fluidity of the brush border membrane. Similarly, altered rates of Na⁺/P_i cotransport occurring during aging and chronic adaptation to a dietary restriction of P_i may also be explained partially by a change of membrane fluidity due to an altered ratio of lipids to cholesterol in the brush border membrane [69, 81].

Metabolic acidosis

Metabolic acidosis provokes an increase of P_i excretion [11, 108]. Transcriptome analysis revealed that NaPi-IIc mRNA is markedly down-regulated in mice after 2 days of metabolic acidosis, less pronounced down-regulation was observed for NaPi-IIa mRNA [86]. In contrast, the abundance of both proteins was increased, consistent with higher Na⁺/P_i cotransport in BBMVs isolated from acidotic mice compared to control animals [87]. Increased urinary P_i excretion was explained by a lower tubular pH-value that decreases the transport rates of NaPi-IIa/c.

Recovery after PTH treatment

In rats injected with a single bolus of PTH, fractional excretion of P_i was maximally increased after 40 min and recovered after 120 minutes. Whereas maximal excretion of P_i correlated with a decrease of the abundance of NaPi-IIa, the levels of NaPi-IIa did not recover parallel to the recovery of P_i excretion [40]. These data indicate that the rate of recovery of the amounts of Na⁺/P_i cotransporters after interventions that provoke their internalization is an important regulatory variable.

Circadian rhythm

Both serum concentration and renal excretion of P_i undergo circadian variations [57]. In rats, fractional excretion of P_i increased substantially between morning and late afternoon, yet, a change in the abundance of NaPi-IIa was not detected [16]. Although part of this observation can be explained by an increase of serum P_i during daytime, it remains to be

shown if fluctuations of the amount of NaPi-IIc and/or Pit-2 may contribute to the diurnal variations of P_i excretion.

Future directions—open questions

Other (Na)/ P_i (co)transporters? Although several Na^+/P_i cotransporters have been localized at the brush border membrane of proximal tubule cells, it is still possible that additional cotransporters may be involved in P_i reabsorption along PT's. If so, it will be of interest to elucidate the relative contribution of each transporter to renal P_i reabsorption in different species. Moreover, the often postulated handling of tubular P_i in distal nephron segments remains to be clarified. Another long-standing open question is the molecular identity of the basolateral exit step of P_i .

Other phosphaturic factors? Recent investigations have added new members to the list of the factors that regulate P_i reabsorption. In the light of a systemic control of renal P_i handling by different axis (bone-kidney, intestine-kidney and eventually others) it is likely that more factors are about to come under the spotlight. Furthermore, the possibility of (a) P_i sensing mechanism(s) (P_i sensor?) remains unanswered.

Control of apical abundance? Although current data suggest that phosphorylation reactions are involved in the control of the apical abundance of Na^+/P_i cotransporters, e.g. by PTH, the signaling cascades of the diverse hormones (including the phosphatonins) that control their apical abundance remain to be elucidated. Moreover, the cellular mechanisms involved in the rapid adaptive responses to altered P_i diets need further investigation.

Trafficking of Na^+/P_i cotransporters Related to the intracellular trafficking of Na/ P_i cotransporters the following questions remain unanswered:

- How are Na^+/P_i cotransporters sorted to the apical membrane?
- Which mechanisms are involved in the routing of Na^+/P_i cotransporters (specifically NaPi-IIa) from the subapical compartment to the lysosomes?

Acknowledgments Work in the authors' laboratories was supported by the Swiss National Science Foundation (SNF), the European Union (EuReGene), the Hartmann Müller Foundation, the Gebert Ruff Foundation, and Transregio SFB 11-Konstanz-Zürich.

References

1. Abraham MI, Woods RE, Breedlove DK, Kempson SA (1992) Renal adaptation to low phosphate diet in diabetic rats. *Am J Physiol* 262:F731–F736
2. Bacic D, Capuano P, Baum M, Zhang J, Stange G, Biber J, Kaissling B, Moe OW, Wagner CA, Murer H (2005) Activation of dopamine D1-like receptors induces acute internalization of the renal Na/phosphate cotransporter NaPi-IIa in mouse kidney and OK cells. *Am J Physiol* 288:F740–F747
3. Bacic D, Hernando N, Traebert M, Lederer E, Völkl H, Biber J, Kaissling B, Murer H (2001) Regulation of the renal type IIa Na/ P_i cotransporter by cGMP. *Pflügers Arch* 443:306–313
4. Bacic D, LeHir M, Biber J, Kaissling B, Murer H, Wagner CA (2006) The renal Na/phosphate cotransporter NaPi-IIa is internalized via the receptor mediated endocytic route in response to parathyroid hormone. *Kidney Int* 69:495–503
5. Bacic D, Schulz N, Biber J, Kaissling B, Murer H, Wagner CA (2003) Involvement of the MAPK-kinase pathway in the PTH-mediated regulation of the proximal tubule type IIa Na/ P_i cotransporter in mouse kidney. *Pflügers Arch* 446:52–60
6. Baum M, Schiavi S, Dwarakanath V, Quigley R (2005) Effect of fibroblast growth factor-23 on phosphate transport in proximal tubules. *Kidney Int* 68:1148–1153
7. Beck L, Karaplis AC, Amizuka N, Hewson AS, Ozawa H, Tenenhouse HS (1998) Targeted inactivation of Npt2 in mice leads to severe renal phosphate wasting, hypercalcuria, and skeletal abnormalities. *Proc Natl Acad Sci (USA)* 95:5372–5377
8. Beers KW, Thompson MA, Chini EN, Dousa TP (1996) beta-Estradiol inhibits Na- P_i cotransport across renal brush border membranes from ovariectomized rats. *Biochem Biophys Res Commun* 221:442–445
9. Bergwitz C, Roslin NM, Tieder M, Loredó-Osti JC, Bastepe M, Abu-Zahra H, Frappier D, Burkett K, Carpenter TO, Anderson D, Garabedian M, Sermet I, Fujiwara TM, Morgan K, Tenenhouse HS, Juppner H (2005) SLC34A3 mutations in patients with hereditary hypophosphatemic rickets with hypercalcuria predict a key role for the sodium-phosphate cotransporter NaPi-IIc in maintaining phosphate homeostasis. *Am J Hum Genet* 78:179–192
10. Berndt TJ, Bielez B, Craig TA, Tebben PJ, Bacic D, Wagner CA, O'Brien S, Schiavi S, Biber J, Murer H, Kumar R (2006) Secreted frizzled-related protein 4 reduces sodium-phosphate cotransporter abundance and activity in proximal tubule cells. *Pflügers Arch* 451:579–587
11. Berndt TJ, Knox FG (1992) Renal regulation of phosphate excretion. In: Seldin DW, Giebisch GH (eds) *The Kidney*, 2nd edn. Williams & Wilkins, Philadelphia, Lippincott, pp 2511–2532
12. Berndt T, Kumar R (2007) Phosphatonins and the regulation of phosphate homeostasis. *Annu Rev Physiol* 69:341–359
13. Berndt T, Thomas LF, Craig TA, Sommer S, Li X, Bergstrahl EJ, Kumar R (2007) Evidence for a signaling axis by which intestinal phosphate rapidly modulates renal phosphate reabsorption. *Proc Natl Acad Sci (USA)* 104:11085–11090
14. Bevington A, Kemp GJ, Graham R, Russell G (1992) Phosphate-sensitive enzymes: A possible molecular basis for cellular disorders of phosphate metabolism. *Clin Chem Enzym Comms* 4:235–257
15. Biber J, Gisler S, Hernando N, Murer H (2005) Protein/protein interactions (PDZ) in proximal tubules. *J Membrane Biol* 203: 111–118
16. Bielez B, Bacic D, Honegger K, Biber J, Murer H, Wagner CA (2006) Unchanged expression of the sodium-dependent phosphate cotransporter NaPi-IIa despite diurnal changes in renal phosphate excretion. *Pflügers Arch* 452:683–689
17. Bielez B, Klaushofer K, Oberbauer R (2004) Renal phosphate loss in hereditary and acquired disorders of bone mineralization. *Bone* 35:1229–1239
18. Bottger P, Hede SE, Grunnet M, Hoyer B, Klaerke DA, Pederson L (2006) Characterization of transport mechanisms and determinants critical for Na-dependent P_i symport of the Pit-family paralogs, human Pit1 and Pit2. *Am J Physiol* 291:C1377–C1387

19. Brautbar N, Walling MW, Coburn JW (1979) Interactions between vitamin D deficiency and phosphorous depletion in the rat. *J Clin Invest* 63:335–341
20. Brown A, Dusso A, Slatopolsky E (2000) Vitamin D. In: Seldin DW, Giebisch GH (eds) *The Kidney*, 3rd edn. Williams & Wilkins, Philadelphia, Lippincott, pp 1047–1090
21. Busch AE, Schuster A, Waldegger S, Wagner CA, Zempel G, Broer S, Biber J, Murer H, Lang F (1996) Expression of a renal type I sodium/phosphate transporter (NaPi-1) induces a conductance in *Xenopus* oocytes permeable for organic and inorganic anions. *Proc Natl Acad Sci (USA)* 83:5347–5351
22. Capuano P, Bacic D, Roos M, Gisler SM, Stange G, Biber J, Kaissling B, Weinman EJ, Shenolikar S, Wagner CA, Murer H (2007) Defective coupling of apical PTH receptors to phospholipase C prevents internalization of the Na-phosphate cotransporter NaPi-1a in NHERF1 deficient mice. *Am J Physiol* 292:C927–C934
23. Capuano P, Bacic D, Stange G, Hernando N, Kaissling B, Pal R, Kocher O, Biber J, Wagner CA, Murer H (2005) Expression and regulation of the renal Na/phosphate cotransporter NaPi-1a in a mouse model deficient for the PDZ protein PDZK1. *Pflügers Arch* 499:392–402
24. Capuano P, Radanovic T, Wagner CA, Bacic D, Kato S, Uchiyama Y, St-Arnaud R, Murer H, Biber J (2005) Intestinal and renal adaptation to a low Pi diet of type Na-Pi cotransporters in vitamin D receptor and 1 α -OHase deficient mice. *Am J Physiol* 288:C429–C434
25. Carpenter T, Ellis BK, Insogna KL, Philbrick WM, Sterpka J, Shinkets R (2005) Fibroblast growth factor 7: An inhibitor of phosphate transport derived from oncogenic osteomalacia-causing tumors. *J Clin Endocrinol Metab* 90:1012–1020
26. Caverzasio J, Bonhour JP (1985) Mechanism of rapid phosphate transport adaptation to a single low Pi meal in rat renal brush border membranes. *Pflügers Arch* 404:227–231
27. Chang Q, Hoefs S, van der Kemp AW, Topala CN, Bindels RJ, Hoenderop JG (2005) The beta-glucuronidase klotho hydrolyzes and activates the TRPV5 channel. *Science* 310:490–493
28. Christensen EI, Nielsen R (2007) Role of megalin and cubulin in renal physiology and pathophysiology. *Rev Physiol Biochem Pharmacol* 158:1–22
29. Collins JF, Ghishan FK (2004) The SLC20 family of proteins: dual functions as sodium-phosphate cotransporters and viral receptors. *Pflügers Arch* 447:647–652
30. Consortium TA (2000) Autosomal dominant hypophosphatemic rickets is associated with mutations in FGF23. *Nat Genet* 26:345–348
31. Cunningham R, Esmaili A, Brown E, Biswas RS, Murtazina R, Donowitz M, Dijkman HB, van der Vlag J, Hogema BM, De Jonge HR, Shenolikar S, Wade JB, Weinman EJ (2008) Urine electrolyte, mineral, and protein excretion in NHERF-2 and NHERF-1 null mice. *Am J Physiol* 294:F1001–F1007
32. Custer M, Lötscher M, Biber J, Murer H, Kaissling B (1994) Expression of Na/Pi cotransport in rat kidney: Localization by RT-PCR and immunohistochemistry. *Am J Physiol* 266:F767–F774
33. Custer M, Meier F, Schlatter E, Greer R, Garcia-Perez A, Biber J, Murer H (1993) Localization of NaPi-1, a NaPi-cotransporter in rabbit kidney proximal tubules. *Pflügers Arch* 424:203–209
34. Deliot N, Hernando N, Liu-Horst Z, Gisler SM, Capuano P, Wagner CA, Bacic D, O'Brien S, Biber J, Murer H (2005) Parathyroid hormone treatment induces dissociation of type IIa Na-Pi cotransporter/ Na/H exchanger regulatory factor-1 complexes. *Am J Physiol* 289:C159–C167
35. Dobbie H, Unwin RJ, Faria NJR, Shirley DG (2008) Matrix extracellular phosphoglycoprotein causes phosphaturia in rats by inhibiting tubular phosphate reabsorption. *Nephrol Dial Transplant* 23:730–733
36. Dousa TP, Kempson SA (1982) Regulation of renal brush border membrane transport of phosphate. *Miner Electrolyte Metab* 7:113–121
37. Farouqi S, Levi M, Soleimani M, Amlal H (2008) Estrogen downregulates the expression of proximal tubule NaPi-1a and causes phosphate wasting and hypophosphatemia. *Kidney Int* 73:1141–1150
38. Forster IC, Köhler K, Biber J, Murer H (2002) Forging the link between structure and function of electrogenic cotransporters: the renal type IIa Na/Pi cotransporter as a case study. *Prog Biophys Mol Biol* 80:69–108
39. Friedlander G (1998) Autocrine/paracrine control of renal phosphate transport. *Kidney Int Suppl* 65:S18–S23
40. Friedlaender MM, Wald H, Dranitzky-Elhalel M, Levi M, Popovtzer MM (2004) Recovery of renal tubule phosphate reabsorption despite reduced levels of sodium-phosphate transporter. *Europ J Endocrin* 151:797–801
41. Gensure RC, Gardella TJ, Jüppner H (2005) Parathyroid hormone and parathyroid hormone related peptide, and their receptors. *Biochim Biophys Acta* 328:666–678
42. Gisler SM, Kittanakom S, Fuster D, Wong V, Bertic M, Radanovic T, Hall RA, Murer H, Biber J, Markovich D, Moe OW, Staglar I (2008) Monitoring protein–protein interactions between the mammalian renal epithelial integral membrane proteins and their PDZ interacting partners using the type II split ubiquitin membrane yeast two-hybrid system. *Molec Cell Proteomics* 7:1362–1377
43. Gisler SM, Pribanic S, Bacic D, Forrer P, Sabourin LA, Tsuji A, Zhao Z, Manser E, Biber J, Murer H (2003) PDZK1: A major scaffold in brush borders of proximal tubular cells. *Kidney Int* 64:1733–1745
44. Gisler SM, Staglar I, Traebert M, Bacic D, Biber J, Murer H (2001) Interaction of the type IIa Na/Pi-cotransporter with PDZ proteins. *J Biol Chem* 276:9206–9213
45. Gmaj P, Murer H (1986) Cellular mechanisms of inorganic phosphate transport in kidney. *Physiol Rev* 66:36–70
46. Guntupalli J, Eby B, Lau K (1982) Mechanism for the phosphaturia of NH₄Cl: dependence on acidemia but not on diet PO₄ or PTH. *Am J Physiol* 242:F552–F560
47. Haggie PM, Stanton BA, Verkman AS (2003) Increased diffusional mobility of CFTR at the plasma membrane after deletion of its C-terminus PDZ binding motif. *J Biol Chem* 279:5494–5500
48. Hammond TG, Yusufi AN, Knox FG, Dousa TP (1985) Administration of atrial natriuretic factor inhibits sodium-coupled transport in proximal tubules. *J Clin Invest* 75:1983–1989
49. Hernando N, Deliot N, Gisler SM, Lederer E, Weinman EJ, Biber J, Murer H (2002) PDZ domain interactions and apical expression of type IIa Na/Pi-cotransporters. *Proc Natl Acad Sci (USA)* 99:11957–11962
50. Hernando N, Forgo J, Biber J, Murer H (2000) PTH-induced downregulation of the type IIa Na/Pi-cotransporter is independent of known endocytic motifs. *J Am Soc Nephrol* 11:1961–1968
51. Hruska KA, Methew S, Lund R, Qui P, Pratt R (2008) Hyperphosphatemia of chronic diseases. *Kidney Int* 74:148–157
52. Hu MC, Zhang J, Shi M, Rosenblatt K, Baum M, Kuro OM, Moe OW (2006) Klotho is a phosphaturic hormone: in vivo evidence. *J Am Soc Nephrol* 17:105a
53. Imura A, Iwano A, Tohyama O, Tsuji Y, Nozaki K, Hashimoto N, Fujimori T, Nabeshima Y (2004) Secreted Klotho protein in sera and CSF: implication for post-translational cleavage in release of Klotho protein from cell membrane. *FEBS Lett* 565: 143–147
54. Ito M, Iidawa S, Izuka M, Haito S, Segawa H, Kuwahata M, Ohkido I, Ohno H, Miyamoto KI (2004) Interaction of a farnesylated protein with renal type IIa Na/Pi cotransporter in

- response to parathyroid hormone and dietary phosphate. *Biochem J* 377:607–616
55. Jüppner H (2007) Novel regulators of phosphate homeostasis and bone metabolism. *Therap Apher Dial* 11:S3–S22
 56. Karim-Jimenez Z, Hernando N, Biber J, Murer H (2000) A dibasic motif involved in parathyroid hormone-induced down-regulation of the type IIa NaPi cotransporter. *Proc Natl Acad Sci (USA)* 97:12896–12901
 57. Kemp GJ, Blumsohn A, Morris BW (1992) Circadian changes in plasma phosphate concentration, urinary phosphate excretion, and cellular phosphate shifts. *Clin Chem* 38:400–402
 58. Kempson SA, Lötscher M, Kaissling B, Biber J, Murer H, Levi M (1995) Parathyroid hormone action on phosphate transporter mRNA and protein in rat renal proximal tubules. *Am J Physiol* 268:F784–F791
 59. Keusch I, Traebert M, Lötscher M, Kaissling B, Murer H, Biber J (1998) Parathyroid hormone and dietary phosphate provoke lysosomal routing of the proximal tubular Na/Pi-cotransporter type II. *Kidney Int* 54:1224–1232
 60. Kido S, Miyamoto K, Mizobuchi H, Taketani Y, Ohkido I, Ogawa N, Kaneko Y, Harashima S, Takeda E (1999) Identification of regulatory sequences and binding proteins in the type II sodium/phosphate cotransporter NPT2 gene responsive to dietary phosphate. *J Biol Chem* 274:28256–28263
 61. Kittler JT, Rostaing P, Schiavo G, Fritschy JM, Olsen R, Triller A, Moss SJ (2001) The subcellular distribution of GABARAP and its ability to interact with NSF suggest a role for this protein in the intracellular transport of GABA(A) receptors. *Mol Cell Neurosci* 18:13–25
 62. Knochel JP (2000) Clinical and physiologic phosphate disturbances. In: Seldin DW, Giebisch GH (eds) *The Kidney*, 3rd edn. Williams & Wilkins, Philadelphia, Lippincott, pp 1905–1934
 63. Köhler K, Forster IC, Lambert G, Biber J, Murer H (2000) The functional unit of the renal type IIa Na⁺/P_i cotransporter is a monomer. *J Biol Chem* 275:26113–26120
 64. Kreusser WJ, Descoedres C, Oda Y, Massry SG, Kurokawa K (1980) Effect of phosphate depletion on renal gluconeogenesis. *Miner Electrolyte Metab* 3:312–323
 65. Kuro OM (2006) Klotho as a regulator of fibroblast growth factor signaling and phosphate/calcium metabolism. *Curr Opin Nephrol Hypertens* 15:437–441
 66. Kurosu H, Ogawa Y, Miyoshi M, Yamamoto M, Nandi A, Rosenblatt KP, Baum MG, Schiavi S, Hu M-C, Moe OW, Kuro-o M (2006) Regulation of fibroblast growth factor-23 signaling by klotho. *J Biol Chem* 281:6120–6123
 67. Lederer ED, Sohi SS, McLeish KR (2000) Parathyroid hormone stimulates extracellular signal-regulated kinase (ERK) activity through two independent signal transduction pathways: role of ERK in sodium-phosphate cotransport. *J Am Soc Nephrol* 11:222–231
 68. Lee SH, Mouradian MM (1999) Up regulation of D1A dopamine receptor gene transcription by estrogen. *Mol Cell Endocrinol* 156:151–157
 69. Levi M, Baird BM, Wilson PV (1990) Cholesterol modulates rat renal brush border membrane phosphate transport. *J Clin Invest* 85:231–237
 70. Levi M, Blaine J, Breusegem S, Takahashi H, Sorribas V, Barry N (2006) Renal phosphate-wasting disorders. *Adv Chron Kidney Dis* 13:155–165
 71. Levi M, Kempson SA, Lötscher M, Biber J, Murer H (1996) Molecular regulation of renal phosphate transport. *J Membrane Biol* 154:1–9
 72. Levi M, Shayman JA, Abe A, Gross SK, McCluer RH, Biber J, Murer H, Lötscher M, Cronin RE (1995) Dexamethasone modulates rat renal brush border membrane phosphate transporter mRNA and protein abundance and glycosylolipid composition. *J Clin Invest* 96:207–216
 73. Levine BS, Ho LD, Pasiecznik K, Coburn JW (1986) Renal adaptation to phosphorous deprivation: characterization of early events. *J Bone Miner Res* 1:33–40
 74. Liu S, Quarles LD (2007) How fibroblast growth factor 23 works. *J Am Soc Nephrol* 18:1637–1647
 75. Lorenz-Depiereux B, Benet-Pages A, Eckstein G, Tenenbaum-Rakover Y, Wagenstaller J, Tiosamo D, Gershoni-Baruch R, Albers N, Lichtner P, Schnabel D, Hochberg Z, Strom TM (2006) Hereditary hypophosphatemic rickets with hypercalcuria is caused by mutations in the sodium-phosphate cotransporter gene SLC34A3. *Am J Hum Genet* 78:193–201
 76. Madjdpour C, Bacic D, Kaissling B, Murer H, Biber J (2004) Segment specific expression of sodium-phosphate cotransporters NaPi IIa and -IIc and interacting proteins in mouse renal proximal tubules. *Pflügers Arch* 448:402–410
 77. Mahon MJ, Cole JA, Lederer ED, Segre GV (2003) Na/H exchanger regulatory factor 1 mediates inhibition of phosphate transport by parathyroid hormone and second messengers by acting at multiple sites in opossum kidney cells. *Mol Endocrinol* 17:2355–2364
 78. McWilliams RR, Breusegem SY, Brodsky KF, Kim E, Levi M, Doctor RB (2005) Shank2E binds NaPi cotransporter at the apical membrane of proximal tubule cells. *Am J Physiol* 289: C1042–C1051
 79. Miyamoto K, Ito M, Tatsumi S, Kuwahata M, Segawa H (2007) New aspect of renal phosphate reabsorption: the type IIc sodium-dependent phosphate transporter. *Am J Nephrol* 27:503–515
 80. Miyamoto KI, Segawa H, Ito M, Kuwahata M (2004) Physiological regulation of renal sodium-phosphate cotransporters. *Japan J Physiol* 54:83–102
 81. Molitoris BA, Alfrey AC, Harris RA, Simon FR (1985) Renal apical membrane cholesterol and fluidity in regulation of phosphate transport. *Am J Physiol* 249:F12–F19
 82. Mühlbauer RC, Bonjour JP, Fleisch H (1981) Tubular handling of Pi: localization of effects of 1,25(OH)₂D₃ and dietary Pi in TPTX rats. *Am J Physiol* 241:F123–F128
 83. Murer H, Forster I, Biber J (2004) The sodium phosphate cotransporter family SLC34. *Pflügers Arch* 447:763–767
 84. Murer H, Hernando N, Forster I, Biber J (2000) Proximal tubular phosphate reabsorption. *Physiol Rev* 80:1373–1409
 85. Nabeshima Y (2002) Klotho: a fundamental regulator of aging. *Ageing Res Rev* 1:627–638
 86. Nowik M, Lecca MR, Velic A, Rehrauer H, Brändli AW, Wagner CA (2008) Genome-wide gene expression profiling reveals renal genes regulated during metabolic acidosis. *Physiol Genomics* 32:322–334
 87. Nowik M, Picard N, Stange G, Capuano P, Tenenhouse HS, Biber J, Murer H (2008) Renal phosphaturia during metabolic acidosis revisited: Molecular mechanisms for decreased renal phosphate reabsorption. *Pflügers Arch* (in press). doi:10.1007/s00424-008-0530-5
 88. Perward F, Zhang MYH, Tenenhouse HS, Portale AA (2007) Fibroblast growth factor 23 impairs phosphorus and vitamin D metabolism in vivo and suppresses 25-hydroxyvitamin D 1α-hydroxylase expression in vitro. *Am J Physiol* 293:F1577–F1583
 89. Pfister MF, Ruf I, Stange G, Ziegler U, Lederer E, Biber J, Murer H (1998) Parathyroid hormone leads to the lysosomal degradation of the renal type II Na⁺/P_i-cotransporter. *Proc Natl Acad Sci (USA)* 95:1909–1914
 90. Picard N, Capuano P, Bacic D, Biber J, Murer H, Le Hir M, Kaissling B, Wagner C (2007) Differential regulation of renal Na/phosphate cotransporters NaPi-IIa and NaPi-IIc in response to parathyroid hormone (PTH). *J Am Soc Nephrol* 18:109A

91. Quarles LD (2003) Evidence for a bone-kidney axis regulating phosphate homeostasis. *J Clin Invest* 112:642–646
92. Radanovic T, Gisler SM, Biber J, Murer H (2007) Topology of the type IIa Na⁺/P_i-cotransporter. *J Membrane Biol* 212:41–49
93. Ravera S, Virkki LV, Murer H, Forster IC (2007) Deciphering PiT transport kinetics and substrate specificity using electrophysiology and flux measurements. *Am J Physiol* 293:C606–C620
94. Reining S, Gisler SM, Biber J, Murer H, Hernando N (2007) Association of NaPi-IIa with the interacting proteins PDZK2 and Gabarap. *J Am Soc Nephrol* 18:123A
95. Rowe PS, de Zoysa PA, Dong R, Wang HR, White KE, Econs MJ, Oudet CL (2000) MEPE, a new gene expressed in bone marrow and tumors causing osteomalacia. *Genomics* 67:54–68
96. Rowe PS, Kumagai Y, Gutierrez G, Garrett IR, Blacher R, Rosen D, Cundy J, Navvab S, Chen D, Drezner MK, Quarles LD, Mundy GR (2004) MEPE has the properties of an osteoblastic phosphatonin and minihibin. *Bone* 34:303–319
97. Saito H, Maeda A, Ohtomo S, Hirata M, Kusano K, Kato S, Ogata E, Segawa H, Miyamoto K, Fukushima N (2005) Circulating FGF-23 is regulated by 1 α , 25-dihydroxyvitamin D3 and phosphorus in vivo. *J Biol Chem* 280:2543–2549
98. Salaun C, Rodrigues P, Heard JM (2001) Transmembrane topology of Pit-2, a phosphate transporter-retrovirus receptor. *J Virol* 75:5584–5592
99. Segawa H, Kaneko I, Takahashi A, Kuwahata M, Ito M, Ohkido I, Tatsumi S, Miyamoto K (2002) Growth-related renal type II Na/Pi cotransporter. *J Biol Chem* 277:19665–19672
100. Segawa H, Kaneko I, Yamanaka S, Ito M, Kuwahata M, Inoue Y, Kato S, Miyamoto K (2004) Intestinal Na–Pi cotransporter adaptation to dietary Pi content in vitamin D receptor null mice. *Am J Physiol* 287:F39–F47
101. Segawa H, Yamanaka S, Ito M, Kuwahata M, Shono M, Yamamoto T, Miyamoto KI (2005) Internalization of renal type IIc Na-Pi cotransporter in response to a high-phosphate diet. *Am J Physiol* 288:F587–F596
102. Segawa H, Yamanaka S, Ohno Y, Onitsuka A, Shiozawa K, Aranami F, Furutani J, Tomoe Y, Ito M, Kuwahata M, Imura A, Nabeshima Y, Miyamoto KI (2007) Correlation between hyperphosphatemia and type II Na-Pi cotransporter activity in *klotho* mice. *Am J Physiol* 292:F769–F779
103. Segawa H, Yamanaka S, Onitsuka A, Tomoe Y, Kuwahata M, Ito M, Taketani Y, Miyamoto KI (2007) Parathyroid hormone-dependent endocytosis of renal type IIc Na-Pi cotransporter. *Am J Physiol* 292:F395–F403
104. Shenolikar S, Voltz JW, Cunningham R, Weinman EJ (2004) Regulation of ion transport by the NHERF family of PDZ proteins. *Physiology* 19:48–54
105. Shenolikar S, Voltz JW, Minkoff CM, Wade J, Weinman EJ (2002) Targeted disruption of the mouse gene encoding a PDZ domain containing protein adaptor, NHERF-1, promotes Npt2 internalization and renal phosphate wasting. *Proc Natl Acad Sci (USA)* 99:11470–11475
106. Shimada T, Hasegawa H, Yamazaki Y, Muto T, Hino R, Takeuchi Y, Fujita T, Nakahara K, Fukumoto S, Yamashita T (2004) FGF-23 is a potent regulator of vitamin D metabolism and phosphate homeostasis. *J Bone Miner Res* 19:429–435
107. Shimada T, Kakitani M, Yamazaki Y, Hasegawa H, Takeuchi Y, Fujita T, Fukumoto S, Tomizuka K, Yamashita T (2004) Targeted ablation of FGF23 demonstrates an essential physiological role of FGF23 in phosphate and vitamin D metabolism. *J Clin Invest* 113:561–568
108. Silve C, Friedlander G (2000) Renal regulation of phosphate excretion. In: Seldin DW, Giebisch GH (eds) *The Kidney*, 3rd edn. Williams & Wilkins, Philadelphia, Lippincott, pp 1885–1903
109. Sommer S, Berndt T, Craig T, Kumar R (2007) The phosphatonins and the regulation of phosphate transport and vitamin D metabolism. *J Steroid Biochem Mole Biol* 103:497–503
110. Spitzer A, Barac-Nieto M (2001) Ontogeny of renal phosphate transport and the process of growth. *Pediatr Nephrol* 16:763–771
111. Syal A, Schiavi S, Chakravarty S, Dwarakanath V, Quigley R, Baum M (2006) Fibroblast growth factor-23 increases mouse PGE2 production in vivo and in vitro. *Am J Physiol* 290:F450–F455
112. Szczepanska-Konkel M, Yusufi AN, VanScoy M, Webster SK, Dousa TP (1986) Phosphonocarboxylic acids as specific inhibitors of Na-dependent transport of phosphate across renal brush border membranes. *J Biol Chem* 261:6375–6383
113. Takeda E, Yamamoto H, Nashiki K, Sato T, Arai H, Taketani Y (2004) Inorganic phosphate homeostasis and the role of dietary phosphorus. *J Cell Mol Med* 8:191–200
114. Taketani Y, Segawa H, Chikamori M, Morita K, Tanaka K, Kido S, Yamamoto H, Iemori Y, Tatsumi S, Tsugawa N, Okano T, Kobayashi T, Miyamoto K, Takeda E (1998) Regulation of type renal Na-dependent inorganic phosphate transporters by 1,25-dihydroxyvitamin D3. Identification of a vitamin D-responsive element in the human NaPi-3 gene. *J Biol Chem* 273:14575–14581
115. Tenenhouse HS, Martel J, Gauthier C, Segawa H, Miyamoto K (2003) Differential effects of Npt2a gene ablation and X-linked Hyp mutation on renal expression of Npt2c. *Am J Physiol* 285:F1271–F1278
116. Tenenhouse HS, Roy S, Martel J, Gauthier C (1998) Differential expression, abundance, and regulation of Na-phosphate cotransporter genes in murine kidney. *Am J Physiol* 275:F527–F534
117. Tenenhouse HS, Sabbagh Y (2002) Novel phosphate-regulating genes in the pathogenesis of renal phosphate wasting disorders. *Pflügers Arch* 444:317–326
118. Tohyama O, Imura A, Iwano A, Freund JN, Henrissat B, Fujimori T, Nabeshima YI (2004) *Klotho* is a novel beta-glucuronidase capable of hydrolyzing steroid beta-glucuronides. *J Biol Chem* 279:9777–9784
119. Traebert M, Volkl H, Biber J, Murer H, Kaisling B (2000) Luminal and contraluminal action of 1–34 and 3–34 PTH peptides on renal type IIa Na/Pi-cotransporter. *Am J Physiol* 278:F792–F798
120. Tröhler U, Bonjour JP, Fleisch H (1976) Renal adaptation to dietary intake in intact and thyroparathyroidectomized rats. *J Clin Invest* 57:264–273
121. Ullrich KJ, Murer H (1982) Sulfate and phosphate transport in rat kidney proximal tubule. *Philos Trans R Soc Lond B Biol Sci* 299:549–558
122. Urakawa I, Yamazaki Y, Shimada T, Iijima K, Hasegawa H, Okawa K, Fujita T, Fukumoto S, Yamashita T (2007) *Klotho* converts canonical FGF receptor into a specific receptor for FGF23. *Nature* 444:770–774
123. Villa-Bellosta R, Barac-Nieto M, Breusegem SY, Barry NP, Levi M, Sorribas V (2008) Interactions of the growth-related, type IIc renal sodium/phosphate cotransporter with PDZ proteins. *Kidney Int* 73:456–464
124. Virkki LV, Biber J, Murer H, Forster IC (2007) Phosphate transporters: a tale of two solute carrier families. *Am J Physiol* 293:F643–F654
125. Virkki LV, Forster IC, Biber J, Murer H (2005) Substrate interactions in the human type IIa sodium-phosphate cotransporter (NaPi-IIa). *Am J Physiol* 288:F969–F981
126. Wagner CA, Hernando N, Forster I, Biber J, Murer H (2008) Genetic defects in renal phosphate handling. In: Genetic diseases of the kidney, edited by Lifton RP, Somlo S, Giebisch G, Seldin DW, Elsevier, (in press).

127. Weber TJ, Liu S, Indridason OS, Quarles LD (2003) Serum FGF23 levels in normal and disordered phosphorus homeostasis. *J Bone Miner Res* 18:1227–1234
128. Weinman EJ, Biswas RS, Peng Q, Shen L, Turner CL, E X, Steplock D, Shenolikar S, Cunningham R (2007) Parathyroid hormone inhibits renal phosphate transport by phosphorylation of serine 77 of sodium-hydrogen exchanger regulatory factor-1. *J Clin Invest* 117:3412–3420
129. Yamashita T, Konishi M, Miyake A, Inui K, Itoh N (2002) Fibroblast growth factor (FGF)-23 inhibits renal phosphate reabsorption by activation of the mitogen-activated protein kinase pathway. *J Biol Chem* 277:28265–28270
130. Yeaman C, Grindstaff KK, Hansen MD, Nelson WJ (1999) Cell polarity: versatile scaffolds keep things in place. *Curr Biol* 15: R515–R517
131. Zajicek HK, Wang H, Puttaparthi K, Halaihel N, Markovich D, Shayman J, Beliveau R, Wilson R, Rogers T, Levi M (2001) Glycosphingolipids modulate renal phosphate transport in potassium deficiency. *Kidney Int* 60: 694–704